

Datasheet for ABIN1859778

anti-CD46 antibody (AA 157-365)






\sim		
Ove	r\/I	Θ VV/
\circ	1 V I	C V V


Overview		
Quantity:	100 μL	
Target:	CD46	
Binding Specificity:	AA 157-365	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CD46 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	MCP (Asp157-Arg365)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against MCP. It has been selected for its ability to recognize MCP in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	
Target Details		
Target:	CD46	
Alternative Name:	Membrane Cofactor Protein (MCP) (CD46 Products)	
Background:	Alternative Names: CD46, MIC10, TLX, TRA2, Trophoblast leukocyte common antigen, CD46	

	Molecule, Complement Regulatory Protein, CD46 Molecule, Complement Regulatory protein
Pathways:	Regulation of Actin Filament Polymerization
Application Details	
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.




Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Mouse Stomach Tissue



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded stomach tissue