

## Datasheet for ABIN1859834

# anti-CCL4 antibody (AA 1-92)

## 1 Image



#### Overview


| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | CCL4   |
| Binding Specificity: | AA 1-92  |
| Reactivity:          | Dog  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This CCL4 antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

#### **Product Details**

| Purpose:          | Polyclonal Antibody to Macrophage Inflammatory Protein 1 Beta (MIP1b)  |
|-------------------|--|
| Immunogen:        | MIP1b (Met1-Asn92)   |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against MIP1b. It has been selected for its ability to recognize MIP1b in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Human  |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |

### Target Details

| Target:                   | CCL4  |
|---------------------------|---|
| Alternative Name:         | MIP1b (CCL4 Products)   |
| Background:               | CCL4, ACT2, G-26, LAG1, MIP1-B, SCYA4, Chemokine C-C-Motif Ligand 4, Small Inducible Cytokine A4, Homologous To Mouse Mip-1b  |
| Application Details       |   |
| Application Notes:        | Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.   |
| Comment:                  | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.   |
| Restrictions:             | For Research Use only   |
| Handling                  |   |
| Format:                   | Liquid  |
| Concentration:            | 500 μg/mL   |
| Buffer:                   | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.   |
| Preservative:             | Sodium azide  |
| Precaution of Use:        | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute |
|                           | azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.   |
| Handling Advice:          |   |
| Handling Advice: Storage: | potentially explosive deposits in lead or copper plumbing.  |
| <u> </u>                  | potentially explosive deposits in lead or copper plumbing.  Avoid repeated freeze-thaw cycles.  |



### **Western Blotting**

Image 1.