

### Datasheet for ABIN1859991

# anti-Neurotensin antibody (AA 21-156)

2 Images




#### Overview

Quantity:	100 μL
Target:	Neurotensin (NTS)
Binding Specificity:	AA 21-156
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Neurotensin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),
	Immunocytochemistry (ICC)
Product Details	
Purpose:	Polyclonal Antibody to Neurotensin (NT)
Immunogen:	RPB203Ra01Recombinant Neurotensin (NT)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NT. It has been selected for its ability
	to recognize NT in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	Neurotensin (NTS)


## **Target Details**

NTS, NTS1, NmN-125, NN, Tail peptide Neuromedin N
Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
20 μg/mL,Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
degradation and precipitation were observed. The loss rate is less than 5% within the expiration
date under appropriate storage condition.
For Research Use only
Liquid
0.5 mg/mL
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
azide-containing compounds in running water before discarding to avoid accumulation of
potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze-thaw cycles.
4 °C,-20 °C
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
detectable loss of activity. Avoid repeated freeze-thaw cycles.
12 months



# Western Blotting

Image 1.



#### **Immunohistochemistry**

**Image 2.** DAB staining on IHC-P; Samples: Rat Kidney Tissue