# antibodies -online.com









# **Images**



| Overview |
|----------|
|          |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | NUMB   |
| Binding Specificity: | AA 376-482   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This NUMB antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

# **Product Details**

| Immunogen:    | NUMB (Leu376-Phe482)   |
|---------------|--|
| Isotype:      | IgG  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against NUMB. It has been selected for its ability to recognize NUMB in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography   |

# **Target Details**

| Target:     | NUMB                            |
|-------------|---------------------------------|
| Abstract:   | NUMB Products                   |
| Background: | Alternative Names: Protein S171 |

| Target Details      |   |
|---------------------|---|
| Pathways:           | Cell-Cell Junction Organization   |
| Application Details |   |
| Application Notes:  | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated  |
|                     | thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious  |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration  |
|                     | date under appropriate storage condition.   |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | Lot specific  |
| Buffer:             | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.   |
| Preservative:       | Sodium azide  |
| Precaution of Use:  | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  |
|                     | Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or   |
|                     | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a   |
|                     | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute   |
|                     | azide-containing compounds in running water before discarding to avoid accumulation of  |
|                     | potentially explosive deposits in lead or copper plumbing.  |

Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Avoid repeated freeze-thaw cycles.

4°C


12 months

Handling Advice:

Storage Comment:


Storage:

Expiry Date:




# **Western Blotting**

Image 1.



# **Western Blotting**

Image 2. Western Blot; Sample: Human A431 cell lysate; Primary Ab: 2μg/ml Rabbit Anti-Human NUMB Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



# **Immunohistochemistry**

**Image 3.** Figure.DAB staining on IHC-P. Samples: Human Tissue