

# Datasheet for ABIN1860121 anti-SERPINB2 antibody (AA 68-313)

# 2 Images



#### Overview

Quantity:	100 μL
Target:	SERPINB2
Binding Specificity:	AA 68-313
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SERPINB2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),
	Immunocytochemistry (ICC)
Product Details	
Purpose:	Polyclonal Antibody to Plasminogen Activator Inhibitor 2 (PAI2)
Immunogen:	RPA531Mu01Recombinant Plasminogen Activator Inhibitor 2 (PAI2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PAI2. It has been selected for its
	ability to recognize PAI2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	SERPINB2


# Target Details

Alternative Name:	PAI2 (SERPINB2 Products)
Target Type:	Amino Acid
Background:	SERPINB2, PLANH2, Serpin Peptidase Inhibitor Clade B Member 2, Monocyte Arg-serpin,
	Placental plasminogen activator inhibitor, Serpin B2, Urokinase inhibitor
Pathways:	Autophagy
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 µg/mL,1:25-100 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	date under appropriate storage condition.  For Research Use only
Restrictions: Handling	
Handling	
	For Research Use only
Handling Format: Concentration:	For Research Use only Liquid
Handling  Format:  Concentration:  Buffer:	For Research Use only  Liquid  0.5 mg/mL
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Handling  Format:  Concentration:  Buffer:  Preservative:	Eiquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
Handling  Format:  Concentration:  Buffer:  Preservative:	For Research Use only  Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
Handling  Format:  Concentration:  Buffer:  Preservative:	Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
Handling  Format:  Concentration:  Buffer:  Preservative:  Precaution of Use:	Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Handling Format:	Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling  Format:  Concentration:  Buffer:  Preservative:  Precaution of Use:  Handling Advice:	Liquid  0.5 mg/mL  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.  Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

### **Images**



## **Western Blotting**

Image 1.



### **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded Liver tissue