antibodies - online.com







anti-PIK3AP1 antibody (AA 397-650)



Images



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 μL	
Target:	PIK3AP1	
Binding Specificity:	AA 397-650	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PIK3AP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	


Product Details

Immunogen:	PIK3AP1 (Glu397-Leu650)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against PIK3AP1. It has been selected for its ability to recognize PIK3AP1 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	

Target Details


Target:	PIK3AP1	
Abstract:	PIK3AP1 Products	
Background:	Alternative Names: BCAP, B-cell adapter for phosphoinositide 3-kinase, B-cell phosphoinositide	

- Target Details		
	3-kinase adapter protein 1	
Pathways:	Activation of Innate immune Response, BCR Signaling, Warburg Effect	
Application Details		
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Human HepG2 cell lysate; Primary Ab: 1μg/ml Rabbit Anti-Rat PIK3AP1 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Rat Tissue