

Datasheet for ABIN1860244 anti-PKM2 antibody (AA 324-531)






Overview

| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | PKM2 |
| Binding Specificity: | AA 324-531 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PKM2 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | PKM2 (Val324-Pro531) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against PKM2. It has been selected for its ability to recognize PKM2 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |
| Target Details | |
| Target: | PKM2 |
| Alternative Name: | Pyruvate Kinase, Muscle (PKM2) (PKM2 Products) |
| Background: | Alternative Names: M2PK, PKM, M2-PK, PKM1, CTHBP, OIP3, PK3, PKM, TCB, THBP1, Pyruvate |
| | |

Target Details

| | kinase muscle isozyme, Thyroid hormone-binding protein 1, Cytosolic thyroid hormone-binding |
|---------------------|---|
| | protein |
| Pathways: | Warburg Effect |
| Application Details | |
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute |
| | azide-containing compounds in running water before discarding to avoid accumulation of |
| | potentially explosive deposits in lead or copper plumbing. |
| Handling Advice: | Avoid repeated freeze-thaw cycles. |
| Storage: | 4 °C |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. |
| Expiry Date: | 12 months |




Western Blotting

Image 1.

Western Blotting

Image 2. Western Blot; Sample: Human HepG2 cell lysate; Primary Ab: 2μg/ml Rabbit Anti-Rat PKM2 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Rat Intestine Tissue