



[Go to Product page](#)

Datasheet for ABIN1860250

## anti-PLA2G2A antibody (AA 22-146)

### 3 Images

#### Overview

Quantity:	100 µL
Target:	PLA2G2A
Binding Specificity:	AA 22-146
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLA2G2A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

#### Product Details

Immunogen:	PLA2G2A (Ser22-Cys146)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PLA2G2A. It has been selected for its ability to recognize PLA2G2A in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

#### Target Details

Target:	PLA2G2A
Alternative Name:	Phospholipase A2, Group IIA (PLA2G2A) ( <a href="#">PLA2G2A Products</a> )
Background:	Alternative Names: PLA2, MOM1, PLA2B, PLA2L, PLA2S, PLAS1, sPLA2(platelets,synovial fluid),

## Target Details

---

Non-pancreatic secretory phospholipase A2, Phosphatidylcholine 2-acylhydrolase 2A

---

Pathways: [Stem Cell Maintenance](#), [Inositol Metabolic Process](#)

---

## Application Details

---

Application Notes: 

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

---

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

---

Restrictions: For Research Use only

---

## Handling

---

Format: Liquid

---

Concentration: Lot specific

---

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

---

Preservative: Sodium azide

---

Precaution of Use: **WARNING:** Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

---

Handling Advice: Avoid repeated freeze-thaw cycles.

---

Storage: 4 °C


---

Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

---


Expiry Date: 12 months

---




### Western Blotting

**Image 1.**



### Immunohistochemistry

**Image 2.** Used in DAB staining on formalin fixed paraffin-embedded testis tissue



### Immunohistochemistry

**Image 3.** DAB staining on IHC-P; Samples: Rat Skin Tissue