

Datasheet for ABIN1860351 anti-COX-1 antibody (AA 23-277)

3 Images



Overview


Quantity:	100 μL
Target:	COX-1
Binding Specificity:	AA 23-277
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COX-1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Cyclooxygenase 1 (COX-1)
Immunogen:	RPA728Ra01Recombinant Cyclooxygenase 1 (COX1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against COX-1. It has been selected for its ability to recognize COX-1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Target:	COX-1
Abstract:	COX-1 Products
Background:	PTGS1, COX3, PCOX1, PGG/HS, PGHS1, PHS1, PTGHS, Prostaglandin Endoperoxide Synthase 1, Prostaglandin H2 Synthase 1, Prostaglandin G/H Synthase And Cyclooxygenase
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	12 months




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Rat Bladder lysate; Primary Ab: 1μg/ml Rabbit Anti-Rat PTGS1 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Rat Liver Tissue