antibodies - online.com







anti-SDF4 antibody (AA 154-362)

Images



\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	SDF4
Binding Specificity:	AA 154-362
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SDF4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details


Immunogen:	SDF4 (Asp154-Phe362)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against SDF4. It has been selected for its ability to recognize SDF4 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	

Target Details

Target:	SDF4	
Abstract:	SDF4 Products	
Background:	Alternative Names: Cab45, RP5-902P8.6, Calcium Binding Protein 45 kDa	

Application Details

1-1-		
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. 	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



Western Blotting

Image 1.

Western Blotting

Image 2. Western Blot; Sample: Mouse Testis lysate; Primary Ab: 1μg/ml Rabbit Anti-Human SDF4 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)