# antibodies - online.com







# anti-SH2D1A antibody (AA 18-115)



# **Images**



( )	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 μL
Target:	SH2D1A
Binding Specificity:	AA 18-115
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH2D1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

### **Product Details**

Immunogen:	SH2D1A (Lys18-Gly115)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against SH2D1A. It has been selected for its ability to recognize SH2D1A in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	

# **Target Details**

Target:	SH2D1A
Alternative Name:	SH2 Domain Containing Protein 1A (SH2D1A) (SH2D1A Products)
Background:	Alternative Names: IMD5, LYP, XLP, MTCP1, DSHP, XLPD, EBVS, SAP, Duncan disease SH2-

Target Details	
	protein, Lymphoproliferative Syndrome, Signaling lymphocytic activation molecule-associated protein
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process
Application Details	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Avoid repeated freeze-thaw cycles.

4°C


12 months

Handling Advice:

Storage Comment:


Storage:

Expiry Date:




## **Western Blotting**

Image 1.



#### **Western Blotting**

Image 2. Western Blot; Sample: Human Jurkat cell lysate; Primary Ab: 2μg/ml Rabbit Anti-Mouse SH2D1A Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



### **Immunohistochemistry**

**Image 3.** Used in DAB staining on fromalin fixed paraffinembedded Lung tissue