antibodies - online.com







anti-TFPI antibody (AA 29-304)

Images



Publication




\sim		
Ove	r\/I	ΔM
\cup	1 V I	CVV

Quantity:	100 μL
Target:	TFPI
Binding Specificity:	AA 29-304
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TFPI antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	TFPI (Asp29-Met304)
	TFPI (Asp29-Met304)
Immunogen:	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its
Immunogen: Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its ability to recognize TFPI in immunohistochemical staining and western blotting.
Immunogen: Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its
Immunogen: Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its ability to recognize TFPI in immunohistochemical staining and western blotting.
Immunogen: Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its ability to recognize TFPI in immunohistochemical staining and western blotting.
Immunogen: Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against TFPI. It has been selected for its ability to recognize TFPI in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography

	pathway inhibitor
Application Details	
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4°C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months
Publications	
Product cited in:	Ranasinghe, Duke, Harvie, McManus: "Kunitz-type protease inhibitor as a vaccine candidate


against schistosomiasis mansoni." in: International journal of infectious diseases: IJID: official publication of the International Society for Infectious Diseases, Vol. 66, pp. 26-32, (2018) (PubMed).

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Human Kidney Tissue