

Datasheet for ABIN1860770 anti-TLR3 antibody (AA 533-705)

2 Images



Overview

Overview	
Quantity:	100 μL
Target:	TLR3
Binding Specificity:	AA 533-705
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TLR3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	TLR3 (Leu533-Leu705)
Immunogen: Isotype:	TLR3 (Leu533-Leu705)
	<u> </u>
Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against TLR3. It has been selected for its
Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against TLR3. It has been selected for its ability to recognize TLR3 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against TLR3. It has been selected for its ability to recognize TLR3 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against TLR3. It has been selected for its ability to recognize TLR3 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography
Isotype: Specificity: Purification: Target Details Target:	IgG The antibody is a rabbit polyclonal antibody raised against TLR3. It has been selected for its ability to recognize TLR3 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography TLR3

Target Details

Pathways:

TLR Signaling, Activation of Innate immune Response, Hepatitis C, Toll-Like Receptors Cascades

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Mouse Tissue