

Datasheet for ABIN1860802
anti-TNFRSF1A antibody (AA 257-458)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	TNFRSF1A
Binding Specificity:	AA 257-458
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFRSF1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Tumor Necrosis Factor Receptor 1 (TNFR1)
Immunogen:	RPB499Ra02Recombinant Tumor Necrosis Factor Receptor 1 (TNFR1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TNFR1. It has been selected for its ability to recognize TNFR1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	TNFRSF1A
Alternative Name:	TNFR1 (TNFRSF1A Products)
Background:	CD120A, P55, TNFRSF1A, TBP1, FPF, TNF-R, TNF-R-I, TNF-R55, TNFAR, TNFR55, TNFR60, P55-R, P60, Tumor necrosis factor receptor 1, Tumor necrosis factor-binding protein 1
Pathways:	NF-kappaB Signaling , Apoptosis , Caspase Cascade in Apoptosis , Hepatitis C , Ubiquitin Proteasome Pathway

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

Handling

detectable loss of activity. Avoid repeated freeze-thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue