antibodies -online.com







anti-WISP2 antibody (AA 15-250)





| \sim | | | | | |
|--------|------|---|------------|----------|---|
| | 1//6 | r | V I | Θ | Λ |


| Overview | | |
|----------------------|--|--|
| Quantity: | 100 μL | |
| Target: | WISP2 | |
| Binding Specificity: | AA 15-250 | |
| Reactivity: | Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This WISP2 antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) | |
| Product Details | | |

| Immunogen: | WISP2 (Leu15-Phe250) | |
|---------------|--|--|
| Isotype: | IgG | |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against WISP2. It has been selected for its ability to recognize WISP2 in immunohistochemical staining and western blotting. | |
| Purification: | Antigen-specific affinity chromatography | |

Target Details


| Target: | WISP2 | |
|-------------|--|--|
| Abstract: | WISP2 Products | |
| Background: | Alternative Names: CCN5, CT58, CTGF-L, CCN family member 5, Connective tissue growth | |

| | factor-like protein, Connective tissue growth factor-related protein 58 | |
|---------------------|--|--|
| Pathways: | WNT Signaling, Growth Factor Binding | |
| Application Dataila | | |
| Application Details | | |
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. | |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated | |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious | |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration | |
| | date under appropriate storage condition. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | Lot specific | |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. | |
| Preservative: | Sodium azide | |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. | |
| | Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or | |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a | |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute | |
| | azide-containing compounds in running water before discarding to avoid accumulation of | |
| | potentially explosive deposits in lead or copper plumbing. | |
| Handling Advice: | Avoid repeated freeze-thaw cycles. | |
| Storage: | 4 °C | |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. | |
| Storage Comment: | Store at 2-0 C for one month. Anquot and store at -50 C for 12 months. | |



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue