

Datasheet for ABIN1862828

anti-Cytokeratin 18 antibody (AA 1-159)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Cytokeratin 18 (KRT18)
Binding Specificity:	AA 1-159
Reactivity:	Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cytokeratin 18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Cytokeratin 18 (CK18)
Immunogen:	RPB231Po01Recombinant Cytokeratin 18 (CK18)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CK18. It has been selected for its ability to recognize CK18 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	Cytokeratin 18 (KRT18)
Alternative Name:	CK18 (KRT18 Products)
Background:	KRT18, CYK18, K18, Keratin 18, Cell proliferation-inducing gene 46 protein, Keratin, type I cytoskeletal 18
Pathways:	Apoptosis, Caspase Cascade in Apoptosis


Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling


Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date: 12 months



Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Human Hela(3u) cell lysate; Primary Ab: 3µg/ml Rabbit Anti-Porcine KRT18 Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)