

Datasheet for ABIN1862981  
**anti-Neuregulin 2 antibody (AA 618-836)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µL
Target:	Neuregulin 2 (NRG2)
Binding Specificity:	AA 618-836
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Neuregulin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## Product Details

Immunogen:	NRG2 (Leu618-Ser836)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NRG2. It has been selected for its ability to recognize NRG2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

## Target Details

Target:	Neuregulin 2 (NRG2)
Alternative Name:	Neuregulin 2 ( <a href="#">NRG2 Products</a> )
Background:	Alternative Names: Don-1, HRG2, NTAK, Neural And Thymus Derived Activator For ErbB

## Target Details

Kinases, Divergent Of Neuregulin-1, Pro-neuregulin-2, membrane-bound isoform

Pathways: [RTK Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#)

## Application Details

Application Notes: 

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: Lot specific

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide


Precaution of Use: WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: 4 °C


Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Expiry Date: 12 months



Immunohistochemistry

**Image 1.** Used in DAB staining on formalin fixed paraffin-embedded stomach tissue



Western Blotting

**Image 2.** Figure. Western Blot; Sample: Recombinant protein.