



[Go to Product page](#)

Datasheet for ABIN1863028
anti-PDIA2 antibody (AA 309-525)

2 Images

Overview

Quantity:	100 µL
Target:	PDIA2
Binding Specificity:	AA 309-525
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDIA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	PDIA2 (Leu309-Leu525)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PDIA2. It has been selected for its ability to recognize PDIA2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	PDIA2
Alternative Name:	Protein Disulfide Isomerase A2 (PDIA2 Products)
Background:	Alternative Names: PDI-A2, PDA2, PDI, PDIR, PDIP, Protein Disulfide Isomerase, Pancreatic,

Target Details

Protein Disulfide Isomerase-Associated 2, Pancreas-specific protein disulfide isomerase

Pathways: [Maintenance of Protein Location](#), [Cell RedoxHomeostasis](#), [Unfolded Protein Response](#)

Application Details

Application Notes:

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide


Precaution of Use: WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: 4 °C


Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Expiry Date: 12 months



Western Blotting

Image 1.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded pancreas tissue