

Datasheet for ABIN1863128

anti-CXCL12 antibody (FITC)



_					
	W	0	rv	10	W

Quantity:	200 μL	
Target:	CXCL12	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CXCL12 antibody is conjugated to FITC	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)	

Product Details

FITC-Linked Polyclonal Antibody to Stromal Cell Derived Factor 1 (SDF1)	
The antibody is a rabbit polyclonal antibody raised against SDF1 conjugated to fitc.	
IgG	
The antibody is a rabbit polyclonal antibody raised against SDF1. It has been selected for its ability to recognize SDF1 in immunohistochemical staining and western blotting.	
Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target:	CXCL12	
Alternative Name:	Stromal Cell Derived Factor 1 (CXCL12 Products)	

Target Details		
Background:	CXCL12, PBSF, P-BSF, SCYB12, SDF1-A, SDF1-B, SDF1A, SDF1B, TLSF-A, TLSF-B, TPAR1, Pre-B	
	Cell Growth-Stimulating Factor, Chemokine C-X-C-Motif Ligand 12	
Pathways:	Regulation of Cell Size, CXCR4-mediated Signaling Events, Negative Regulation of intrinsic	
	apoptotic Signaling	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100	
	Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by	
	end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze/thaw cycles	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	