# antibodies -online.com







# anti-NPTX2 antibody

**Images** 



#### Overview

Quantity:	200 μL
Target:	NPTX2
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This NPTX2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**


Immunogen:	The antibody is a mouse monoclonal antibody raised against NPTX2.
Isotype:	IgG
Specificity:	It has been selected for its ability to recognize NPTX2 in immunohistochemical staining and Western blotting.
Purification:	Affinity Chromatography

#### **Target Details**

Target:	NPTX2
Alternative Name:	Neuronal Pentraxin II (NPTX2 Products)


### **Application Details**

Application Notes:	Western blotting: 1:100-400
	Immunocytochemistry in formalin fixed cells: 1:100-500
	Immunohistochemistry in formalin fixed frozen section: 1:100-500
	Immunohistochemistry in paraffin section: 1:50-200
	Enzyme-linked Immunosorbent Assay: 1:100-200
	Optimal working dilutions must be determined by end user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	Supplied as solution form in PBS, pH7.4, containing 0.02 % NaN3, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze/thaw cycles
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months



## Western Blotting

Image 1.



#### **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded Pancreas tissue