antibodies - online.com







anti-CUTA antibody (AA 33-179)

Images



\sim					
	1//6	r	V I	Θ	Λ

Background:

Quantity:	100 μL
Target:	CUTA
raiget.	COTA
Binding Specificity:	AA 33-179
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CUTA antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	ACHAP (Arg33-Pro179)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ACHAP. It has been selected for its
	ability to recognize ACHAP in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	CUTA
Alternative Name:	Acetylcholinesterase Associated Protein (CUTA Products)

Alternative Names: CUTA, C6orf82, CutA Divalent Cation Tolerance Homolog, Brain

	acetylchol	linesterase	putative	membrane	anchor
--	------------	-------------	----------	----------	--------

Application Details

• Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Application Notes: Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in

paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated


thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration

date under appropriate storage condition.

Restrictions: For Research Use only


Handling

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue