# antibodies - online.com







# anti-AP2M1 antibody (AA 170-435)





$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	AP2M1
Binding Specificity:	AA 170-435
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AP2M1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

#### **Product Details**


Immunogen:	AP2m1 (Arg170-Cys435)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against AP2m1. It has been selected for its ability to recognize AP2m1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

### **Target Details**

Target:	AP2M1
Alternative Name:	Adaptor Related Protein Complex 2 Mu 1 (AP2M1 Products)
Background:	Alternative Names: AP50, CLAPM1, Clathrin-Associated/Assembly/Adaptor Protein,Medium 1,


## Target Details

	Plasma Membrane Adaptor AP-2 50kDA Protein, Clathrin Coat Adaptor Protein AP50
Pathways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, EGFR Downregulation, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Format:  Concentration:	Liquid  Lot specific
Concentration:	Lot specific
Concentration: Buffer:	Lot specific  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Concentration:  Buffer:  Preservative:	Lot specific  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Concentration:  Buffer:  Preservative:  Precaution of Use:	Lot specific  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Concentration:  Buffer:  Preservative:  Precaution of Use:  Handling Advice:	Lot specific  PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.  Sodium azide  WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.  Avoid repeated freeze-thaw cycles.



#### **Western Blotting**

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



#### **Immunohistochemistry**

**Image 2.** Figure.DAB staining on IHC-P. Samples: Human Tissue