antibodies - online.com







anti-APPBP2 antibody (AA 396-585)

Images



Overview

Quantity:	100 μL
Target:	APPBP2
Binding Specificity:	AA 396-585
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APPBP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	APPBP2 (Lys396-Cys585)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against APPBP2. It has been selected for its ability to recognize APPBP2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	APPBP2	
Alternative Name: Amyloid Beta Precursor Protein Binding Protein 2 (APPBP2 Products)		
Background:	Alternative Names: PAT1, Protein Interacting With APP Tail 1,Cytoplasmic Tail, Amyloid protein-	

bindina	protein 2	. Protein	interacting	with	APP tail 1

Application Details

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling


Comment:

Format:	Liquid		
Concentration:	Lot specific		
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.		
Preservative:	Sodium azide		
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.		
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or		
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a		
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute		
	azide-containing compounds in running water before discarding to avoid accumulation of		
	potentially explosive deposits in lead or copper plumbing.		
Handling Advice:	Avoid repeated freeze-thaw cycles.		
Storage:	4 °C		
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.		
Expiry Date:	12 months		



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue