antibodies - online.com







anti-ARHGDIB antibody (AA 2-201)





_					
U	V	er	V	Ie	W

Quantity:	100 μL	
Target:	ARHGDIB	
Binding Specificity:	AA 2-201	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ARHGDIB antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	

Product Details

Immunogen:	ARHGDIb (Thr2-Glu201)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against ARHGDIb. It has been selected for its ability to recognize ARHGDIb in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	

Target Details

Target:	ARHGDIB	
Abstract:	ARHGDIB Products	
Background:	Alternative Names: D4, GDIA2, GDID4, LYGDI, Ly-GDI, RAP1GN1, RhoGDI2	

Pathways:

Caspase Cascade in Apoptosis

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	



Immunohistochemistry

Image 1. Figure.DAB staining on IHC-P. Samples: Human Tissue



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.