

Datasheet for ABIN1866875

anti-Butyrylcholinesterase antibody (AA 345-570)

2 Images



Overview

Quantity:	100 μL
Target:	Butyrylcholinesterase (BCHE)
Binding Specificity:	AA 345-570
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Butyrylcholinesterase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Butyrylcholinesterase (BCHE)
Immunogen:	RPC348Ra01Recombinant Butyrylcholinesterase (BCHE)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against BCHE. It has been selected for its ability to recognize BCHE in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Target:	Butyrylcholinesterase (BCHE)
Alternative Name:	BCHE (BCHE Products)
Background:	BuChE, CHE1, E1, Pseudocholinesterase, Cholinesterase, Acylcholine acylhydrolase,
	Butyrylcholine esterase, Choline esterase II
Pathways:	Peptide Hormone Metabolism
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	The state of the s
	date under appropriate storage condition.
Restrictions:	For Research Use only
Restrictions: Handling	
Handling	For Research Use only
Handling Format:	For Research Use only Liquid
Handling Format: Concentration:	For Research Use only Liquid 0.5 mg/mL
Handling Format: Concentration: Buffer:	For Research Use only Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Handling Format: Concentration: Buffer: Preservative:	For Research Use only Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Handling Format: Concentration: Buffer: Preservative:	For Research Use only Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Handling Format: Concentration: Buffer: Preservative:	For Research Use only Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
Handling Format: Concentration: Buffer: Preservative:	For Research Use only Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
Handling Format: Concentration: Buffer: Preservative:	Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
Handling Format: Concentration: Buffer: Preservative:	Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Handling Format: Concentration: Buffer: Preservative: Precaution of Use:	Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Format: Concentration: Buffer: Preservative: Precaution of Use: Handling Advice:	Liquid 0.5 mg/mL PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. ProClin, Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing. Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Rat Kidney Tissue