

### Datasheet for ABIN1866905

# anti-Biliverdin Reductase antibody (AA 1-125)






### Overview

Overview	
Quantity:	100 μL
Target:	Biliverdin Reductase (BLVRA)
Binding Specificity:	AA 1-125
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Biliverdin Reductase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	BLVRA (Met1-His125)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against BLVRA. It has been selected for its ability to recognize BLVRA in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	Biliverdin Reductase (BLVRA)
Alternative Name:	Biliverdin Reductase A (BLVRA Products)


## **Application Details**

<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>
The thermal stability is described by the loss rate. The loss rate was determined by accelerated
thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious
degradation and precipitation were observed. The loss rate is less than 5% within the expiration
date under appropriate storage condition.
For Research Use only
Liquid
Lot specific
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
azide-containing compounds in running water before discarding to avoid accumulation of
potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze-thaw cycles.
4 °C
Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
12 months




### **Western Blotting**

Image 1. Western Blot; Sample: Human Lung lysate; Primary Ab: 1μg/ml Rabbit Anti-Human BLVRA Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



#### **Western Blotting**

**Image 2.** Figure. Western Blot; Sample: Recombinant protein.



### **Immunohistochemistry**

**Image 3.** Used in DAB staining on fromalin fixed paraffinembedded liver tissue