antibodies -online.com





anti-CRIP1 antibody (AA 2-77)

2 Images



Go to Product page

Overview

| Overview | |
|----------------------|---|
| Quantity: | 100 μL |
| Target: | CRIP1 |
| Binding Specificity: | AA 2-77 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CRIP1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | CRIP1 (Pro2-Lys77) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against CRIP1. It has been selected for its |

Target Details

Purification:

| Target: | CRIP1 |
|-------------------|--|
| Alternative Name: | Cysteine Rich Protein 1, Intestinal (CRIP1 Products) |
| Background: | Alternative Names: CRP1, CRHP, CRIP, Cysteine-rich heart protein, Cysteine-rich intestinal |


Antigen-specific affinity chromatography

ability to recognize CRIP1 in immunohistochemical staining and western blotting.

Expiry Date:


12 months

| l arget Details | |
|--|--|
| | protein |
| Application Details | |
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| | |
| Concentration: | Lot specific |
| Concentration: Buffer: | Lot specific PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| | |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Buffer: Preservative: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of |
| Buffer: Preservative: Precaution of Use: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Sodium azide WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing. |



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue