antibodies -online.com





anti-CST5 antibody (AA 21-142)

3 Images




Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Overview	
Quantity:	100 μL
Target:	CST5
Binding Specificity:	AA 21-142
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CST5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	CST5 (Gly21-Val142)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CST5. It has been selected for its ability to recognize CST5 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	CST5
Alternative Name:	Cystatin 5 (CST5) (CST5 Products)
Background:	Alternative Names: Cystatin D


Application Details

Application Details		
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiratio date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	e: Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	




Western Blotting

Image 1.



Western Blotting

Image 2. Western Blot; Sample: Human Saliva; Primary Ab: 5μg/ml Rabbit Anti-Human CST5 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue