# antibodies -online.com







## anti-ECF antibody (AA 24-97)





$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	ECF	
Binding Specificity:	AA 24-97	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ECF antibody is un-conjugated	
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	ECF (Gly24-Pro97)	
9	26. (6.)2657)	
	lgG	
Isotype:		
Isotype:	IgG	
Isotype: Specificity: Purification:	lgG  The antibody is a rabbit polyclonal antibody raised against ECF. It has been selected for its	
Isotype: Specificity: Purification:	IgG  The antibody is a rabbit polyclonal antibody raised against ECF. It has been selected for its ability to recognize ECF in immunohistochemical staining and western blotting.	
Isotype: Specificity: Purification: Target Details	IgG  The antibody is a rabbit polyclonal antibody raised against ECF. It has been selected for its ability to recognize ECF in immunohistochemical staining and western blotting.	
Isotype: Specificity:	IgG  The antibody is a rabbit polyclonal antibody raised against ECF. It has been selected for its ability to recognize ECF in immunohistochemical staining and western blotting.  Antigen-specific affinity chromatography	

C-Motif Ligand 11, Small Inducible C	tokine Subfamily A(	Cys-Cys	)Member 11
--------------------------------------	---------------------	---------	------------

## **Application Details**

#### Application Notes:

• Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

#### Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### Restrictions:

For Research Use only


## Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months



## **Western Blotting**

Image 1.



### **Immunohistochemistry**

**Image 2.** Figure.DAB staining on IHC-P. Samples: Human Tissue