



[Go to Product page](#)

Datasheet for ABIN1868507
anti-IGF2BP3 antibody (AA 2-343)

3 Images

Overview

Quantity:	100 µL
Target:	IGF2BP3
Binding Specificity:	AA 2-343
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGF2BP3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	IGF2BP3 (Asn2-Ile343)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IGF2BP3. It has been selected for its ability to recognize IGF2BP3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	IGF2BP3
Alternative Name:	Insulin Like Growth Factor 2 mRNA Binding Protein 3 (IGF2BP3 Products)
Background:	Alternative Names: MP-3, IMP3, KOC1, VICKZ3, CT98, IGF II mRNA Binding Protein 3,

Target Details


Cancer/Testis Antigen 98, KH domain-containing protein overexpressed in cancer

Application Details

Application Notes:	<ul style="list-style-type: none">Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months




Immunohistochemistry

Image 1. DAB staining on IHC-P;;Samples: Human Kidney Tissue



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded pancreas tissue



Western Blotting

Image 3. Figure. Western Blot; Sample: Recombinant protein.