

Datasheet for ABIN1868520 anti-IGFBP3 antibody (AA 140-271)

2 Images



Overview

Quantity:	100 μL
Target:	IGFBP3
Binding Specificity:	AA 140-271
Reactivity:	Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGFBP3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),
	Immunocytochemistry (ICC)
Product Details	
Purpose:	Polyclonal Antibody to Insulin Like Growth Factor Binding Protein 3 (IGFBP3)
Immunogen:	RPA054Po01Recombinant Insulin Like Growth Factor Binding Protein 3 (IGFBP3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IGFBP3. It has been selected for its
	ability to recognize IGFBP3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	IGFBP3

Target Details

Alternative Name:	IGFBP3 (IGFBP3 Products)
Background:	BP53, IBP3, Growth Hormone-Dependent Binding Protein, Acid Stable Subunit Of The 140 K IGF Complex
Pathways:	Myometrial Relaxation and Contraction, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development, Regulation of Carbohydrate Metabolic Process, Autophagy, Smooth Muscle Cell Migration, Growth Factor Binding
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions: Handling	For Research Use only
Handling	For Research Use only
Handling Format:	For Research Use only Liquid
Handling Format: Concentration:	For Research Use only Liquid 500 µg/mL
Handling Format: Concentration: Buffer:	For Research Use only Liquid 500 μg/mL PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Handling Format: Concentration: Buffer: Preservative:	Eiquid 500 μg/mL PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300. Dithiothreitol (DTT), ProClin WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of
Handling Format: Concentration: Buffer: Preservative: Precaution of Use:	Eiquid 500 μg/mL PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300. Dithiothreitol (DTT), ProClin WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:


12 months

Images



Western Blotting

Image 1. Western Blot; Sample: Porcine Serum; ;Primary Ab: 1μg/ml Rabbit Anti-Porcine IGFBP3 Antibody;Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody;(Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.