# antibodies - online.com







# anti-MIA antibody (AA 36-127)

**Images** 



$\sim$	
( )\/\	rview
$\cup$	

Quantity:	100 μL
Target:	MIA
Binding Specificity:	AA 36-127
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MIA antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC)

# **Product Details**


Immunogen:	MIA1 (Ala36-Phe127)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MIA1. It has been selected for its ability to recognize MIA1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

# **Target Details**

Target:	MIA
Alternative Name:	Melanoma Inhibitory Activity Protein 1 (MIA1) (MIA Products)
Background:	Alternative Names: CD-RAP, Melanoma-derived growth regulatory protein


# **Application Details**

10 10 10 10 10 10 10 10 10 10 10 10 10 1	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.  Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	4 °C
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
Expiry Date:	12 months




# **Western Blotting**

Image 1.



### **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded stomach tissue



# **Immunohistochemistry**

**Image 3.** DAB staining on IHC-P; Samples: Rat Stomach Tissue