antibodies - online.com





anti-MTR antibody (AA 923-1265)

Images



\sim						
	1//	Д	r۱	/1	\triangle	٨

O V CI V I C V V	
Quantity:	100 μL
Target:	MTR
Binding Specificity:	AA 923-1265
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
FIUUUUL DELAIIS	


Immunogen:	MTR (Ser923-Asp1265)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MTR. It has been selected for its ability to recognize MTR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	MTR
Alternative Name:	5-Methyltetrahydrofolate Homocysteine Methyltransferase (MTR Products)
Background:	Alternative Names: CbIG, MS, MeSe, MetH Methionine Synthase, Vitamin-B12 dependent


Target Details

ethionine synthase ethionine Biosynthetic Process
ethionine Biosynthetic Process
Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
ne thermal stability is described by the loss rate. The loss rate was determined by accelerated ermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious egradation and precipitation were observed. The loss rate is less than 5% within the expiration at under appropriate storage condition.
or Research Use only
quid
ot specific
3S, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
odium azide
ARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Void contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or ve contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a hysician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute tide-containing compounds in running water before discarding to avoid accumulation of otentially explosive deposits in lead or copper plumbing.
void repeated freeze-thaw cycles.
°C
ore at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
2 months



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue