antibodies - online.com







anti-NR1D2 antibody (AA 249-578)

Images



_		
()Ver	view	

Quantity:	100 μL
Target:	NR1D2
Binding Specificity:	AA 249-578
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR1D2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	


Product Details

Immunogen:	NR1D2 (Lys249-Pro578)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NR1D2. It has been selected for its ability to recognize NR1D2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details


Target:	NR1D2
Alternative Name:	Nuclear Receptor Subfamily 1, Group D, Member 2 (NR1D2 Products)
Background:	Alternative Names: HZF2, BD73, EAR-1r, RVR, Rev-ErbA Beta, Orphan nuclear hormone receptor

BD73, Rev-erb alpha-related receptor, V-erbA-related protein 1-related
Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway
 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
For Research Use only
Liquid
Lot specific
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze-thaw cycles.
4 °C
Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
12 months



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Rat Tissue