antibodies -online.com





anti-PCDHB16 antibody (AA 35-347)

2 Images



Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Background:


Quantity:	100 μL
Target:	PCDHB16
Binding Specificity:	AA 35-347
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	PCDHb16 (Val35-Val347)
	PCDHb16 (Val35-Val347) IgG
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against PCDHb16. It has been selected for
Immunogen: Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against PCDHb16. It has been selected for its ability to recognize PCDHb16 in immunohistochemical staining and western blotting.
Immunogen: Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against PCDHb16. It has been selected for its ability to recognize PCDHb16 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography
Immunogen: Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against PCDHb16. It has been selected for its ability to recognize PCDHb16 in immunohistochemical staining and western blotting.

PCDHbeta 16

Alternative Names: PCDHB8a, PCDH3X, PCDH-BETA16, ME1, Cadherin ME1, Protocadherin-3x,


Application Details

Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. 				
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated				
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious				
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.				
Restrictions:	For Research Use only				
Handling					
Format:	Liquid				
Concentration:	Lot specific				
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.				
Preservative:	Sodium azide				
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.				
Handling Advice:	Avoid repeated freeze-thaw cycles.				
Storage:	4 °C				
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.				
Expiry Date:	12 months				



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue