

Datasheet for ABIN1870061 anti-PTGDS antibody (AA 1-199)

2 Images




Go to Product page

\sim			
	ve	r\/	٨
\cup	V C	1 V I	٧V

Quantity:	100 μL		
Target:	PTGDS		
Binding Specificity:	AA 1-199		
Reactivity:	Human		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This PTGDS antibody is un-conjugated		
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),		
	Immunocytochemistry (ICC)		
Product Details			
Purpose:	Polyclonal Antibody to Prostaglandin D2 Synthase (PTGDS)		
Immunogen:	RPB640Hu01Recombinant Prostaglandin D2 Synthase (PTGDS)		
Isotype:	IgG		
Specificity:	The antibody is a rabbit polyclonal antibody raised against PTGDS. It has been selected for its		
	ability to recognize PTGDS in immunohistochemical staining and western blotting.		
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography		
Target Details			
Target:	PTGDS		


Target Details

Alternative Name:	PTGDS (PTGDS Products)	
Background:	HPGDS, GSTS, H-PGDS, PDS, PGD2, PGDS, PGDS2, Glutathione S-Transferase Sigma,	
	Glutathione-requiring prostaglandin D synthase, Prostaglandin D2 Synthase, Hematopoietic	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	



Western Blotting

Image 1.



Immunohistochemistry

Image 2. DAB staining on IHC-P;;Samples: Human Stomach cancer Tissue