

Datasheet for ABIN1870127
anti-PTPRG antibody (AA 848-1119)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	PTPRG
Binding Specificity:	AA 848-1119
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPRG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	PTPRG (Phe848-Ala1119)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PTPRG. It has been selected for its ability to recognize PTPRG in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	PTPRG
Alternative Name:	Protein Tyrosine Phosphatase Receptor Type G (PTPRG Products)
Background:	Alternative Names: PTPR-G, HPTPG, PTPG, R-PTP-GAMMA, RPTPG, Receptor-type tyrosine-

Target Details

protein phosphatase gamma

Application Details

Application Notes: • Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: Lot specific

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide


Precaution of Use: WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Handling Advice: Avoid repeated freeze-thaw cycles.

Storage: 4 °C


Storage Comment: Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Expiry Date: 12 months



Immunohistochemistry

Image 1. Figure.DAB staining on IHC-P. Samples: Human Tissue



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.