# antibodies -online.com







## anti-SIAE antibody (FITC)



#### Overview

Quantity:	200 μL
Target:	SIAE
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SIAE antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

#### **Product Details**

1 Toddet Details	
Immunogen:	The antibody is a rabbit polyclonal antibody raised against SIAE conjugated to fitc.
Isotype:	IgG
Specificity:	It has been selected for its ability to recognize SIAE in immunohistochemical staining and Western blotting.
Purification:	Affinity Chromatography
Target Details	

Target:	SIAE
Alternative Name:	Sialic Acid Acetylesterase (SIAE Products)

### **Application Details**

Western blotting: 1:100-400
Immunocytochemistry in formalin fixed cells: 1:100-500
Immunohistochemistry in formalin fixed frozen section: 1:100-500
Immunohistochemistry in paraffin section: 1:50-200
Enzyme-linked Immunosorbent Assay: 1:100-200
Optimal working dilutions must be determined by end user.
For Research Use only
Liquid
Lot specific
Supplied as solution form in PBS, pH7.4, containing 0.02 % NaN3, 50 % glycerol.
Sodium azide
WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
azide-containing compounds in running water before discarding to avoid accumulation of
potentially explosive deposits in lead or copper plumbing.
Avoid repeated freeze/thaw cycles
4 °C
Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.
12 months