antibodies -online.com





anti-SOCS2 antibody (AA 1-198)



Images



Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μL
Target:	SOCS2
Binding Specificity:	AA 1-198
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	SOCS2 (Met1-Val198)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against SOCS2. It has been selected for i ability to recognize SOCS2 in immunohistochemical staining and western blotting.	
Purification: Antigen-specific affinity chromatography		

Target Details

Target:	SOCS2
Alternative Name: Suppressors Of Cytokine Signaling 2 (SOCS2) (SOCS2 Products)	
Background:	Alternative Names: CIS2, Cish2, SOCS-2, SSI-2, SSI2, STATI2, Cytokine-inducible SH2 protein 2, STAT-induced STAT inhibitor 2

JAK-STAT Signaling, Response to Growth Hormone Stimulus

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


Handling

Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.	
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or	
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a	
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute	
	azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	




Western Blotting

Image 1. Western Blot; Sample: Rat Liver lysate; Primary Ab: 1μg/ml Rabbit Anti-Rat SOCS2 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Figure.DAB staining on IHC-P. Samples: Rat Tissue