antibodies -online.com







anti-SYCN antibody (AA 38-134)

2 Images



Go to Product page

| _ | | | | | | |
|---|-----|---|---------------|----|---|----|
| | 1// | Д | $r \setminus$ | /1 | Р | ۱۸ |


Background:

| Quantity: | 100 μL |
|----------------------|--|
| Target: | SYCN |
| Binding Specificity: | AA 38-134 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SYCN antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | SYCN (Cys38-Ser134) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against SYCN. It has been selected for its ability to recognize SYCN in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |
| Target Details | |
| Target: | SYCN |
| Alternative Name: | Syncollin (SYCN Products) |

Alternative Names: SYL, INSSA1, Insulin Synthesis Associated 1


Application Details

| 10 10 10 10 10 10 10 10 10 10 10 10 10 1 | | | | |
|--|---|--|--|--|
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. | | | |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated | | | |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious | | | |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration | | | |
| | date under appropriate storage condition. | | | |
| Restrictions: | For Research Use only | | | |
| Handling | | | | |
| Format: | Liquid | | | |
| Concentration: | Lot specific | | | |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. | | | |
| Preservative: | Sodium azide | | | |
| Precaution of Use: | WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or | | | |
| | eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a | | | |
| | physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute | | | |
| | azide-containing compounds in running water before discarding to avoid accumulation of | | | |
| | potentially explosive deposits in lead or copper plumbing. | | | |
| Handling Advice: | Avoid repeated freeze-thaw cycles. | | | |
| Storage: | 4 °C | | | |
| Storage Comment: | Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months. | | | |
| Expiry Date: | 12 months | | | |
| | | | | |



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Human Tissue