antibodies - online.com







anti-Transferrin Receptor 2 antibody (AA 493-785)

Images



_						
()	V	ρ	rv	16	$\supset I$	Λ

Quantity:	100 μL
Target:	Transferrin Receptor 2 (TFR2)
Binding Specificity:	AA 493-785
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Transferrin Receptor 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	

roduct Details

Immunogen:	TFR2 (His493-Asn785)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TFR2. It has been selected for its ability to recognize TFR2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	Transferrin Receptor 2 (TFR2)	
Alternative Name:	Transferrin Receptor 2 (TFR2 Products)	
Background:	Alternative Names: HFE3, TFRC2	

Target Details	
Pathways:	Transition Metal Ion Homeostasis
Application Details	
Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or


	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of	
	potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze-thaw cycles.	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	
Expiry Date:	12 months	

eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a




Western Blotting

Image 1. Western Blot; Sample: Rat Liver lysate; Primary Ab: 2μg/ml Rabbit Anti-Rat TFR2 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded liver tissue