antibodies -online.com







anti-TRIP6 antibody (AA 281-471)

Images



\sim	
Over\	$/1 \triangle 1 \Lambda /$
OVEIV	/ I C V V

Quantity:	100 μL
Target:	TRIP6
Binding Specificity:	AA 281-471
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIP6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	TRIP6 (Gly281-Glu471)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TRIP6. It has been selected for its ability to recognize TRIP6 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TRIP6
Alternative Name:	Thyroid Hormone Receptor Interactor 6 (TRIP6 Products)
Background:	Alternative Names: OIP1, ZRP-1, Opa-interacting protein 1, Zyxin-related protein 1

Target Details	
Pathways:	Cell-Cell Junction Organization
Application Details	
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled.
	Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or
	eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a
	physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute
	azide-containing compounds in running water before discarding to avoid accumulation of
	potentially explosive deposits in lead or copper plumbing.

Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.

Avoid repeated freeze-thaw cycles.

4°C


12 months

Handling Advice:

Storage Comment:


Storage:

Expiry Date:



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Figure.DAB staining on IHC-P. Samples: Mouse Tissue