# antibodies - online.com







# anti-Angiostatin antibody (FITC)



#### Overview

Quantity:	200 μL
Target:	Angiostatin (ANG)
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Angiostatin antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

#### **Product Details**

Immunogen:	The antibody is a rabbit polyclonal antibody raised against ANG conjugated to fitc.
Isotype:	IgG
Specificity:	It has been selected for its ability to recognize ANG in immunohistochemical staining and Western blotting.
Purification:	Affinity Chromatography

## Target Details

Target:	Angiostatin (ANG)
Alternative Name:	Angiostatin (ANG Products)
Pathways:	Lipid Metabolism

## **Application Details**

Expiry Date:

12 months

A Prince of the Control of the Contr		
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Reconstitution:	Reconstitute in sterile PBS, pH 7.2 - pH 7.4.	
Concentration:	Lot specific	
Buffer:	Supplied as solution form in PBS, pH7.4, containing 0.02 % NaN3, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	WARNING: Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.	
Handling Advice:	Avoid repeated freeze/thaw cycles	
Storage:	4 °C	
Storage Comment:	Store at 2-8 °C for one month. Aliquot and store at -80 °C for 12 months.	