

Datasheet for ABIN1881047
anti-ADPRH antibody (C-Term)[Go to Product page](#)

1 Image

4 Publications

Overview

Quantity:	400 µL
Target:	ADPRH
Binding Specificity:	AA 306-334, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADPRH antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This ADPRH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 306-334 amino acids from the C-terminal region of human ADPRH.
Clone:	RB42002
Isotype:	Ig Fraction
Predicted Reactivity:	B
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	ADPRH
Alternative Name:	ADPRH (ADPRH Products)

Target Details

Background: The enzyme encoded by this gene catalyzes removal of mono-ADP-ribose from arginine residues of proteins in the ADP-ribosylation cycle. Unlike the rat and mouse enzymes, which require DTT for maximal activity, the human enzyme is DTT-independent.

Molecular Weight: 39507

NCBI Accession: [NP_001116](#)

UniProt: [P54922](#)

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications

Product cited in: Hyrskyluoto, Bruelle, Lundh, Do, Kivinen, Rappou, Reijonen, Waltimo, Petersén, Lindholm, Korhonen: "Ubiquitin-specific protease-14 reduces cellular aggregates and protects against mutant huntingtin-induced cell degeneration: involvement of the proteasome and ER stress-activated kinase IRE1?." in: **Human molecular genetics**, Vol. 23, Issue 22, pp. 5928-39, (2014) ([PubMed](#)).


Davila, Froeling, Tan, Bonnard, Boland, Snippe, Hibberd, Seielstad: "New genetic associations detected in a host response study to hepatitis B vaccine." in: **Genes and immunity**, Vol. 11, Issue 3, pp. 232-8, (2010) ([PubMed](#)).

Chen, Qin, Li, Walters, Wilson, Mei, Wilson: "The proteasome-associated deubiquitinating enzyme Usp14 is essential for the maintenance of synaptic ubiquitin levels and the development of neuromuscular junctions." in: **The Journal of neuroscience : the official journal of the Society for Neuroscience**, Vol. 29, Issue 35, pp. 10909-19, (2009) ([PubMed](#)).

Nagai, Kadowaki, Maruyama, Takeda, Nishitoh, Ichijo: "USP14 inhibits ER-associated degradation via interaction with IRE1alpha." in: **Biochemical and biophysical research communications**, Vol. 379, Issue 4, pp. 995-1000, (2009) ([PubMed](#)).

Mines, Goodwin, Limbird, Cui, Fan: "Deubiquitination of CXCR4 by USP14 is critical for both CXCL12-induced CXCR4 degradation and chemotaxis but not ERK activation." in: **The Journal of biological chemistry**, Vol. 284, Issue 9, pp. 5742-52, (2009) ([PubMed](#)).

Images



Western Blotting

Image 1. ADPRH Antibody (C-term) (ABIN1881047 and ABIN2838918) western blot analysis in HeLa cell line lysates (35 µg/lane). This demonstrates the ADPRH antibody detected the ADPRH protein (arrow).