


Datasheet for ABIN1881099

anti-BAD antibody (pSer32)





Go to Product page

| Overview | |
|----------------------|--|
| Quantity: | 400 μL |
| Target: | BAD |
| Binding Specificity: | pSer32 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BAD antibody is un-conjugated |
| Application: | Dot Blot (DB) |
| Product Details | |
| Immunogen: | This BAD Antibody is generated from rabbits immunized with a KLH conjugated synthetic |
| | phosphopeptide corresponding to amino acid residues surrounding S32 of human BAD. |
| Clone: | RB41775 |
| Isotype: | lg Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Target Dataila | |
| Target Details | |
| Target: | BAD |
| Alternative Name: | BAD (BAD Products) |
| Background: | The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are |

Target Details

known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq].

Molecular Weight:

18392

NCBI Accession:

NP_004313, NP_116784

UniProt:

Q92934

Pathways:

MAPK Signaling, PI3K-Akt Signaling, RTK Signaling, Apoptosis, Fc-epsilon Receptor Signaling
Pathway, Positive Regulation of Peptide Hormone Secretion, Carbohydrate Homeostasis,
Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process,
Hepatitis C, CXCR4-mediated Signaling Events

Application Details

Application Notes:

DB: 1:500

Restrictions:

For Research Use only

Handling

| Format: |
|---------|
| Dff |

Liquid

Buffer:

Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative:

Sodium azide

Precaution of Use:

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

should be handled by trained staff only.

Storage:

4 °C,-20 °C

Expiry Date:

6 months

Publications

Product cited in:

Galmiche, Ezzoukhry, François, Louandre, Sabbagh, Nguyen-Khac, Descamps, Trouillet, Godin, Regimbeau, Joly, Barbare, Duverlie, Mazière, Chatelain: "BAD, a proapoptotic member of the BCL2 family, is a potential therapeutic target in hepatocellular carcinoma." in: **Molecular cancer**

research: MCR, Vol. 8, Issue 8, pp. 1116-25, (2010) (PubMed).

Yu, Sun, Shen, Gao, Fan, Sun: "Expression of the apoptosis-related genes BCL-2 and BAD in human breast carcinoma and their associated relationship with chemosensitivity." in: **Journal of experimental & clinical cancer research : CR**, Vol. 29, pp. 107, (2010) (PubMed).

Images



Dot Blot

Image 1. Dot blot analysis of BAD Antibody (Phospho S32) Phospho-specific Pab r on nitrocellulose membrane. 50 ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6 μg per ml.