ANTIBODIES ONLINE

Datasheet for ABIN1881102 anti-BAT1 antibody (C-Term)

4 Images

1 Publication



Overview

| Quantity: | 400 µL |
|----------------------|---|
| Target: | BAT1 (DDX39) |
| Binding Specificity: | AA 351-380, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This BAT1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

Product Details

| Immunogen: | This BAT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 351-380 amino acids from the C-terminal region of human BAT1. |
|-----------------------|--|
| Clone: | RB22515 |
| Isotype: | Ig Fraction |
| Predicted Reactivity: | M, Rat, B, C, D, Pig |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

Target:

BAT1 (DDX39)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN1881102 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|----------------------------------|---|
| Alternative Name: | BAT1 (DDX39 Products) |
| Background: | Component of the THO subcomplex of the TREX complex. The TREX complex specifically associates with spliced mRNA and not with unspliced pre-mRNA. It is recruited to spliced mRNAs by a transcription-independent mechanism. Binds to mRNA upstream of the exon- junction complex (EJC) and is recruited in a splicing-and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export. The recruitment occurs via an interaction between THOC4 and the cap-binding protein NCBP1. UAP56 functions as a bridge between THOC4 and the THO complex. The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production. The recruitment of the TREX complex to the intronless viral mRNA occurs via an interaction between KSHV ORF57 protein and THOC4. Splice factor that is required for the first ATP-dependent step in spliceosome assembly and for the interaction of U2 snRNP with the branchpoint. It has both RNA-stimulated ATP binding/hydrolysis activity and ATP-dependent RNA unwinding activity. Even with the stimulation of RNA, the ATPase activity is weak. It can only hydrolyze ATP but not other NTPs. The RNA stimulation of ATPase activity does not have a strong preference for the sequence and length of the RNA. However, ssRNA stimulates the ATPase activity much more strongly than dsRNA. It can unwind 5' or 3' overhangs or blunt end RNA duplexes in vitro. The ATPase and helicase activities are not influenced by U2AF2 and THOC4. |
| Molecular Weight: | 48991 |
| NCBI Accession: | NP_004631, NP_542165 |
| UniProt: | Q13838 |
| Pathways: Application Details | Ribonucleoprotein Complex Subunit Organization |
| Application Notes: | WB: 1:1000, WB: 1:1000, IHC-P: 1:10~50, FC: 1:10~50 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide. |
| Preservative: | Sodium azide |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN1881102 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

| Н | land | ling |
|---|------|------|
| | | 9 |

| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
|--------------------|--|
| Storage: | 4 °C,-20 °C |
| Expiry Date: | 6 months |
| Publications | |
| Product cited in: | Choudhary, Kumar, Gnad, Nielsen, Rehman, Walther, Olsen, Mann: "Lysine acetylation targets |
| | protein complexes and co-regulates major cellular functions." in: Science (New York, N.Y.), Vol. |
| | 325, Issue 5942, pp. 834-40, (2009) (PubMed). |

Images





Image 1. Formalin-fixed and paraffin-embedded human kidney reacted with BAT1 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry, clinical relevance has not been evaluated.



Western Blotting

Image 2. BAT1 Antibody (C-term) (ABIN1881102 and ABIN2842459) western blot analysis in A431,Hela,Jurkat cell line lysates (35 µg/lane).This demonstrates the BAT1 antibody detected the BAT1 protein (arrow).

Images



Western Blotting

Image 3. All lanes : Anti-BAT1 Antibody (C-term) at 1:1000 dilution Lane 1: Hela whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.

Please check the product details page for more images. Overall 4 images are available for ABIN1881102.