



[Go to Product page](#)

Datasheet for ABIN1881223

anti-COPG antibody (C-Term)

1 Image

2 Publications

Overview

| | |
|----------------------|-------------------------------------|
| Quantity: | 400 µL |
| Target: | COPG |
| Binding Specificity: | AA 721-750, C-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This COPG antibody is un-conjugated |
| Application: | Western Blotting (WB) |

Product Details

| | |
|---------------|--|
| Immunogen: | This COPG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 721-750 amino acids from the C-terminal region of human COPG. |
| Clone: | RB41787 |
| Isotype: | Ig Fraction |
| Purification: | This antibody is purified through a protein A column, followed by peptide affinity purification. |

Target Details

| | |
|-------------------|--|
| Target: | COPG |
| Alternative Name: | COPG (COPG Products) |
| Background: | The coatamer is a cytosolic protein complex that binds to dilysine motifs and reversibly |

Target Details

associates with Golgi non-clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. Coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins, the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors (By similarity).

Molecular Weight: 97718

NCBI Accession: [NP_057212](#)

UniProt: [Q9Y678](#)

Application Details

Application Notes: WB: 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Expiry Date: 6 months

Publications

Product cited in: Dai, Liu, Liu, Zhang, Wang, Jin, Qian, Wang, Zhao, Wu, Xiong, Chang, Sun, Yang, Hoffman, Liu: "Anti-metastatic Efficacy of Traditional Chinese Medicine (TCM) Ginsenoside Conjugated to a VEGFR-3 Antibody on Human Gastric Cancer in an Orthotopic Mouse Model." in: **Anticancer research**, Vol. 37, Issue 3, pp. 979-986, (2017) ([PubMed](#)).

Irrthum, Karkkainen, Devriendt, Alitalo, Vikkula: "Congenital hereditary lymphedema caused by a


mutation that inactivates VEGFR3 tyrosine kinase." in: **American journal of human genetics**, Vol. 67, Issue 2, pp. 295-301, (2000) ([PubMed](#)).

Galland, Karamysheva, Pebusque, Borg, Rottapel, Dubreuil, Rosnet, Birnbaum: "The FLT4 gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor." in: **Oncogene**, Vol. 8, Issue 5, pp. 1233-40, (1993) ([PubMed](#)).

Pajusola, Aprelikova, Korhonen, Kaipainen, Pertovaara, Alitalo, Alitalo: "FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines." in: **Cancer research**, Vol. 52, Issue 20, pp. 5738-43, (1992) ([PubMed](#)).

Galland, Karamysheva, Mattei, Rosnet, Marchetto, Birnbaum: "Chromosomal localization of FLT4, a novel receptor-type tyrosine kinase gene." in: **Genomics**, Vol. 13, Issue 2, pp. 475-8, (1992) ([PubMed](#)).

Images



Western Blotting

Image 1. All lanes : Anti-COPG Antibody (C-term) at 1:1000 dilution Lane 1: whole cell lysate Lane 2: K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 98 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.