

Datasheet for ABIN1881290

anti-EIF3E antibody (AA 248-276)**2** Images**5** Publications[Go to Product page](#)

Overview

Quantity:	400 µL
Target:	EIF3E
Binding Specificity:	AA 248-276
Reactivity:	Human, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF3E antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	This EIF3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-276 amino acids from the Central region of human EIF3E.
Clone:	RB40515
Isotype:	Ig Fraction
Predicted Reactivity:	X, B, C, Pr, M, Rat
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	EIF3E
Alternative Name:	EIF3E (EIF3E Products)

Target Details

Background:	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA ⁱ and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay (NMD), may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.
Molecular Weight:	52221
NCBI Accession:	NP_001559
UniProt:	P60228
Pathways:	Ribonucleoprotein Complex Subunit Organization , Hepatitis C

Application Details

Application Notes:	WB: 1:1000. WB: 1:1000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Expiry Date:	6 months

Publications

Product cited in:	Grzmil, Rzymiski, Milani, Harris, Capper, Saunders, Salhan, Ragoussis, Norbury: "An oncogenic role of eIF3e/INT6 in human breast cancer." in: Oncogene , Vol. 29, Issue 28, pp. 4080-9, (2010) (
-------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

[PubMed](#)).


Zhou, Sandercock, Fraser, Ridlova, Stephens, Schenauer, Yokoi-Fong, Barsky, Leary, Hershey, Doudna, Robinson: "Mass spectrometry reveals modularity and a complete subunit interaction map of the eukaryotic translation factor eIF3." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 105, Issue 47, pp. 18139-44, (2008) ([PubMed](#)).

Morris, Wittmann, Jäck, Jalinot: "Human INT6/eIF3e is required for nonsense-mediated mRNA decay." in: **EMBO reports**, Vol. 8, Issue 6, pp. 596-602, (2007) ([PubMed](#)).

Masutani, Sonenberg, Yokoyama, Imataka: "Reconstitution reveals the functional core of mammalian eIF3." in: **The EMBO journal**, Vol. 26, Issue 14, pp. 3373-83, (2007) ([PubMed](#)).


Sirchia, Luparello: "Mid-region parathyroid hormone-related protein (PTHrP) and gene expression of MDA-MB231 breast cancer cells." in: **Biological chemistry**, Vol. 388, Issue 5, pp. 457-65, (2007) ([PubMed](#)).

Images



Western Blotting

Image 1. EIF3E Antibody (Center) (ABIN1881290 and ABIN2838389) western blot analysis in K562 cell line lysates (35 µg/lane). This demonstrates the EIF3E antibody detected the EIF3E protein (arrow).



Western Blotting

Image 2. (DANRE) eif3eb Antibody (Center) (ABIN1881290 and ABIN2838389) western blot analysis in zebra fish brain tissue lysates (35 µg/lane). This demonstrates the (DANRE) eif3eb antibody detected the (DANRE) eif3eb protein (arrow).