



[Go to Product page](#)

Datasheet for ABIN1881326  
**anti-ETV7 antibody (C-Term)**

1 Image

### Overview

Quantity:	400 µL
Target:	ETV7
Binding Specificity:	AA 313-341, C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ETV7 antibody is un-conjugated
Application:	Western Blotting (WB)

### Product Details

Immunogen:	This ETV7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 313-341 amino acids from the C-terminal region of human ETV7.
Clone:	RB42837
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

### Target Details

Target:	ETV7
Alternative Name:	ETV7 ( <a href="#">ETV7 Products</a> )
Background:	Transcriptional repressor, binds to the DNA sequence 5'-CCGGAAGT-3'. Isoform A does not

## Target Details

seem to have a repressor activity. Isoform C does not seem to have a repressor activity.

Molecular Weight: 38998

NCBI Accession: [NP\\_001193964](#), [NP\\_001193965](#), [NP\\_001193966](#), [NP\\_001193967](#), [NP\\_001193968](#),  
[NP\\_001193969](#), [NP\\_001193970](#), [NP\\_057219](#)

UniProt: [Q9Y603](#)

## Application Details

Application Notes: WB: 1:2000

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Expiry Date: 6 months

## Images



### Western Blotting

**Image 1.** All lanes : Anti-ETV7 Antibody (C-term) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: K562 whole cell lysate Lane 4: NCI- whole cell lysate Lane 5: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5 % NFDM/TBST.