



[Go to Product page](#)

Datasheet for ABIN1881337
anti-FAP antibody (N-Term)

3 Images

3 Publications

Overview

Quantity:	400 µL
Target:	FAP
Binding Specificity:	AA 15-41, N-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	This Mouse Fap antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-41 amino acids from the N-terminal region of mouse Fap.
Clone:	RB40867
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

Target Details

Target:	FAP
Alternative Name:	Fap (FAP Products)
Background:	In association with DPP4 is involved in the pericellular proteolysis of the extracellular matrix

Target Details

(ECM), the migration and invasion of endothelial cells into the ECM (By similarity). May have a role in tissue remodeling during development and wound healing, and contribute to invasiveness in malignant cancers.

Molecular Weight: 87945

NCBI Accession: [NP_032012](#)

UniProt: [P97321](#)

Pathways: [Tube Formation](#)

Application Details

Application Notes: WB: 1:2000. WB: 1:500. IHC-P: 1:100

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Expiry Date: 6 months

Publications

Product cited in: Sun, Sun, Chen, Liao, He, Wang, Chen, Hu, Qiu: "microRNA-27b shuttled by mesenchymal stem cell-derived exosomes prevents sepsis by targeting JMJD3 and downregulating NF-κB signaling pathway." in: **Stem cell research & therapy**, Vol. 12, Issue 1, pp. 14, (2021) ([PubMed](#)).


Reithmair, Buschmann, Märte, Kirchner, Hagl, Kaufmann, Pfob, Chouker, Steinlein, Pfaffl, Schelling: "Cellular and extracellular miRNAs are blood-compartment-specific diagnostic targets in sepsis." in: **Journal of cellular and molecular medicine**, Vol. 21, Issue 10, pp. 2403-2411, (2018) ([PubMed](#)).

Youn, Friesen, Kishimoto, Henne, Kurat, Ye, Ceccarelli, Sicheri, Kohlwein, McMahon, Andrews: "Dissecting BAR domain function in the yeast Amphiphysins Rvs161 and Rvs167 during endocytosis." in: **Molecular biology of the cell**, Vol. 21, Issue 17, pp. 3054-69, (2010) ([PubMed](#)).

Qian, Shi, Pang, Wu, Yu, Li, Wang, Zhou: "[Identification and expression of two new secretory proteins associated with prostate cancer]." in: **Yi chuan = Hereditas / Zhongguo yi chuan xue hui bian ji**, Vol. 32, Issue 3, pp. 235-41, (2010) ([PubMed](#)).


Hwangbo, Kim, Lee, Lee: "Activation of the integrin effector kinase focal adhesion kinase in cancer cells is regulated by crosstalk between protein kinase Calpha and the PDZ adapter protein mda-9/Syntenin." in: **Cancer research**, Vol. 70, Issue 4, pp. 1645-55, (2010) ([PubMed](#)).

Images




Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemical analysis of (ABIN1881337 and ABIN2838726) on paraffin-embedded Mouse kidney tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH 9.0). Samples were incubated with primary Antibody (1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Western Blotting

Image 2. Anti-Mouse Fap Antibody (N-term) at 1:2000 dilution + mouse kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 88 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.



Western Blotting

Image 3. All lanes : Anti-Mouse Fap Antibody (N-term) at 1:500 dilution Lane 1: Mouse kidney lysate Lane 2: Mouse ovary lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 88 kDa Blocking/Dilution buffer: 5 % NFDm/TBST.